MEDITECH Expanse TIP SHEET





Pharmacy

MAR Pain score outside of PRN range

Medication Start Stop Statt Percocet 5/325 mg 1 tab PO Q4H PRN Generic: axycODONE/accetamiophen 03/31/25 11:14 Active Dispense: 1 Tablet/1 tab Give: 1 Tablet (1 tab total) 03/31/25 11:14 Active PRN Reason PRN Reason PRN Reason PRN Reason	e	<u>eMAR</u> Example #1 Documented verbal numeric pain score is HIGHER than the PRN range on the order.
Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source Source	Werk 31 11:16 by 8RU9401 Percocet 5/325 mg 1 tab PO Q4H PRN • Yets No • Yets No • Verbal numeric • • Worop-laker FACEs pain rating scale (Non-verbal) • • CPOT critical care Pain Observation Tool (Non-communicative) • • FLACC Pediatire Face, Leg, Activity, Crying, Consolability (Non-communicative) • • NPASS Neonatal Pain, Agitation, and Sedation Scale (Non-communicative) • • • No pain • • • • Nid pain • • • • Nid pain • • • • Nid pain • • • • • Nid pain • • • • • • • • • • • • • • • • • • • • •	
Warning Image: This medication has been orders are: pain scale 4-6. The pain in This pain score is HIGHER than override reason to continue. * Override Comment is required Override Reason to continue. * Override Comment is required Override Reason to continue. * Override Sea Potent Pt PrefLess Intrusion	Nurse will be presented with an over-ride reason why they are using a less potent medication or a less intrusive route of administration. The response will be saved on the Rx audit of the medication.	

MAR Pain score outside of PRN range

Admin Trowsneet Provinger Monograph Links	Example #2
Medication Start Stop Status	
Percocet 5/325 mg 1 tab PO Q4H PRN Generic: oxyCODONE/acetaminophen	Documented verbal numeric
Dispense: 1 Tablet/1 tab 03/31/25 11:14 Active Give: 1 Tablet (1 tab total) PRIK Passen	pain score is LOWER than
pain scale 4-6	the PRN range on the order.
Source Percocet 5/325 mg 1 tab PO Q4H PRN	401
Assessments	
Pain Assessment MAR Assessing for Pain Medication Administering/assessing for pain management O Yes O No	
Vain Scale Utilized Pain scale utilized Pain scale utilized O Verbal numeric O Wong Baker FACES pain rating scale (Non-verbal)	
 ○ Wong-Baker FACES pain rating scale (Non-verbal) ○ CPDT Critical Care Pain Observation Tool (Non-communicative) ○ FLACC Pediatric Face, Leg, Activity, Crying, Consolability (Non-communicative) ○ NIFS Neonatal/Infant Pain Scale (Non-communicative) 	
(N-PASS Neonatal Pain, Agitation, and Sedation Scale (Non-communicative)	
Numeric Pain Scale Numeric Pain Scale Numeric pain scale O 0 - No pain O 2 - Mild pain O 4 - Moderate pain O 5 - Moderate pain O 7 - Severe pai	iain 08 - Sever
Vong-Baker Pain Scale Vong-Baker Pain Scale COT	
> FLACC > NIPS	
NPASS Van Intensity Pain intensity Z	
rum maany *	
Error	This will present a LIADD
	This will present a HARD
	stop to nursing, and they
	should choose an
	appropriate medication
	for the pain score.
This medication has been ordered for pain. The parameters for administration are: pain scale 4-6. The pain intensity score is: 2	
This pain score is LOWER than the PRN pain score range, use a pain medication with an appropriate PRN range.	
with an appropriate PKN range.	

