

# MEDITECH Expanse TIP SHEET


## Pharmacy/OM


### Cervical Ripening Cervidil Protocol

#### Provider Order Entry

Cervical Ripening indication will default. Provider will need to enter a criteria for use.


Protocol: Cervical Ripening Cervidil

\* Criteria for Cervidil Use: 

Rx Indication:  Cervical Ripening

- If checklist cannot be completed and all items are not documented in chart, oxytocin or cervical ripening should not be initiated.
- 1. Indication for induction/augmentation/cervical ripening documented in chart
- 2. Pelvis is documented by provider to be clinically adequate for vaginal delivery
- 3. Estimated fetal weight is less than 5000 grams in non-diabetic patient (4500 grams for diabetic)
- 4. Gestational age documented
- 5. Provider with C-Section privileges is aware of the induction and readily available

Protocol: Cervical Ripening Cervidil

\* Criteria for Cervidil Use: 

Rx Indication:

Fetal Growth Rest. (<10%)  
Oligohydramnios  
Prolonged Preg. (>42 wks)  
Other

- If checklist cannot be completed and all items are not documented in chart, oxytocin or cervical ripening should not be initiated.
- 1. Indication for induction/augmentation/cervical ripening documented in chart
- 2. Pelvis is documented by provider to be clinically adequate for vaginal delivery
- 3. Estimated fetal weight is less than 5000 grams in non-diabetic patient (4500 grams for diabetic)
- 4. Gestational age documented
- 5. Provider with C-Section privileges is aware of the induction and readily available

If one of the three criteria are not applicable, provider will enter Other.

Protocol

Cervical Ripening Cervidil

\* Criteria for Cervidil Use:

- Other
- If cervidil exceptions are not met, use alternative options including single mechanical methods (foley, cook), misoprostol or combination methods.

\* Other Reason for Cervidil Use:

Rx Indication: Cervical Ripening

- If checklist cannot be completed and all items are not documented in chart, oxytocin or cervical ripening should not be initiated.

If Other is chosen, guidance text will display, and provider is prompted to enter other reason for use.

## Pharmacy Verification

1 of 8 Selected

Orders with Activity

Instructions Status Source Select Patient

View Order Data

Protocol

Protocol Queries

Criteria for Cervidil Use: Other

If cervidil exceptions are not met, use alternative options including single mechanical methods (foley, cook), misoprostol or combination methods.

Other Reason for Cervidil Use: free text reason

Rx Indication: Cervical Ripening

If checklist cannot be completed and all items are not documented in chart, oxytocin or cervical ripening should not be initiated.

1. Indication for induction/augmentation/cervical ripening documented in chart

2. Pelvis is documented by provider to be clinically adequate for vaginal delivery

3. Estimated fetal weight is less than 5000 grams in non-diabetic patient (4500 grams for diabetic)

4. Gestational age documented

5. Provider with C-Section privileges is aware of the induction and readily available

6. Status of cervix is assessed and documented, including all elements of Bishop score

7. Repression is assessed and documented as applicable

Medication Taper Protocol Chemo Comments Instructions Queries Screenings Provider

Protocol

Criteria for Cervidil Use:

- Other
- If cervidil exceptions are not met, use alternative options including single mechanical methods (foley, cook), misoprostol or combination methods.

Other Reason for Cervidil Use: free text reason

Rx Indication: IN.CERVRIIP

1 of 3 Goto 2

Cancel Save

Medication			
<b>Cervidil, 10 mg (See Protocol) VAGINAL ONCE ONE</b>	Start	Stop	Status
Generic: dinoprostone	07/03/25 08:45	07/03/25 08:46	Active
Dispense: 1 Vaginal Suppository/10 mg Give: 1 Vaginal Suppository (10 mg total)			
<a href="#">Detail</a> <a href="#">History</a> <a href="#">Flowsheet</a> <a href="#">Monograph</a> <a href="#">AssocData</a> <a href="#">Prot/Taper</a> <a href="#">Order</a> <a href="#">Links</a>			
Protocol: Cervical Ripening Cervidil			
<b>Criteria for Cervidil Use:</b> Other • If cervidil exceptions are not met, use alternative options including single mechanical methods (foley, cook), misoprostol or combination methods. <b>Other Reason for Cervidil Use:</b> free text reason <b>Rx Indication:</b> Cervical Ripening			
• If checklist cannot be completed and all items are not documented in chart, oxytocin or cervical ripening should not be initiated. • 1. Indication for induction/augmentation/cervical ripening documented in chart • 2. Pelvis is documented by provider to be clinically adequate for vaginal delivery • 3. Estimated fetal weight is less than 5000 grams in non-diabetic patient (4500 grams for diabetic) • 4. Gestational age documented • 5. Provider with C-Section privileges is aware of the induction and readily available • 6. Status of cervix is assessed and documented, including all elements of Bishop score • 7. Presentation is assessed and documented as cephalic			
All items listed are documented in the patient chart: Yes			
If no, why:			

eMAR

# Expanse TIP SHEET

## Discharge Plan – Forms added to Signature Page



When the inpatient discharge plan is printed it will now display “Discharge Plan” and “Discharge Medication List” on the Forms section of what the patient received. “Discharge Medication List” will appear only if the medications have been finalized

Discharge Signature Page		Page: 1 of 1
		Date: 08/07/25 12:54
Frisbie Memorial Hospital		
test,dcplan 50 M 01/01/75		MR# F000546185 Acct# A00000206199 Adm Date 04/09/25
<u>The above-named patient and/or representative has received the following:</u>		
Patient Education CAP A Healthier You		
Forms	Discharge Plan	
	Discharge Medication List	
	Care Assure	
	Suicide Prevention Resources	
I have read and understand the instructions given to me by my caregivers.		
test,dcplan		
Print Patient Name		
Patient (or Guardian) Signature		Date/Time
Caregiver/RN/Doctor Signature		Date/Time
Printed on 08/07/25 at 12:54		

Expanse TIP SHEET

Discharge Plan – Medication Strength Display in Discharge Packet



<div><div>Discharge Medications</div><div>Finalized</div><div><div>amoxicillin-pot clavulanate [Augmentin] 500-125 mg tablet 1 tab</div><div>PO BID Qty: 10 ORF</div><div>Rx</div></div><div><div>ciprofloxacin HCl 0.3 % drops 1 drp ophthalmic (eye) Q4H Qty: 5 ORF</div><div>Rx</div></div><div><div>Rx Instructions: 1 drp into the eye(s);</div><div>diphenhydramine-acetaminophen [Acetaminophen PM] 25-500 mg tablet 1 tab PO BEDTIME PRN (Reason: sedation) Qty: 10 ORF</div><div>Rx</div></div><div><div>docusate sodium [Colace] 100 mg capsule 100 mg PO DAILY Qty: 10 ORF</div><div>Rx</div></div><div><div>insulin detemir U-100 100 unit/mL solution 10 unit subcut BEDTIME Qty: 10 ORF</div><div>Rx</div></div><div><div>oxycodone-acetaminophen [Percocet] 5-325 mg tablet 1 tab</div><div>PO Q8H PRN (Reason: pain) Qty: 10 ORF</div><div>Rx</div></div><div><div>vancomycin 1,000 mg recon soln 480 mg IV BID Qty: 10 ORF</div><div>Rx</div></div></div>
--

The medication strength that currently displays on the discharge plan will now be displayed on the inpatient discharge medication home list when the discharge packet is printed

# MEDITECH Expanse TIP SHEET

## Bariatric Surgery – Diet Alert

Conflicts

TRAINING,OCSSPatient10 43 F 01/11/1982 Allergy/Adv: chocolate, Penicillins

(More ▾)

Close

▼ Regular Diet

\*Override

\*Erase

SCREENING TYPE	DESCRIPTION	SEVERITY
▼ Order Rule	OM BariatricPt Ord Conflict	
Rule Message	This patient account indicates the patient has a history of bariatric surgery. Please ensure the appropriate diet order has been selected for this patient.	

If a patient has a history of bariatric surgery, the system will now alert ordering users if they attempt to submit a diet order that is not a **Bariatric** or **NPO** diet.

This rule will trigger for the following diet order selections:

- Regular Diet
- Low Sodium Diet
- Low Fat Diet
- Clear Liquid Diet
- Consistent Carb/Diabetic Diet
- Dysphagia/IDDSI Diet
- Fiber Restricted Diet
- Full Liquid Diet
- Pediatric Diet
- Renal Diet
- Adult Tube Feeding
- Pediatric Tube Feeding

enicillins

(More ▾)

Close

\*Override

\*Erase

SEVERITY

history of bariatric surgery. Please ensure the patient.

Users have the option to:

- Select **Override** to continue with the selected order.
- Select **Erase** to remove the order from the order queue.

# CHG0397870– Add rule to Discharge w Parameters order to require a selection for Vital Signs

Changes go in effect 08/20/25

- Per CR 1495 – Require the temperature range, systolic blood pressure range, diastolic blood pressure range, heart rate range, respiration range, and O2 saturation range queries when “Vital Signs” group response is selected from the Discharge Parameter query.

- If any of the temperature range, systolic blood pressure range, diastolic blood pressure range, heart rate range, respiration range, and O2 saturation are filled out the other queries become unrequired.

# MEDITECH Expanse TIP SHEET

## Provider Consult – Edit Restriction

A work-around was recently discovered in which users were able to Edit existing Non-Telehealth Provider Consult orders and enter a provider group into the “Consulting Provider” standard field. This is prevented during initial order entry via an order conflict rule. However, no such guardrail exists within the order edit routine.

This workaround was causing massive numbers of physicians to be related to the patient and causing numerous downstream issues.

Order Summary

SAVE 1

CURRENT ENTER RECONCILE

Next Req Field ▶

ORDERS DATE STATUS X ALL

Consult to Hospitalist Routine 07/23/25 15:27 \* Edited X

STOP DATE

Routine \* Edited 07/23/25 15:27

Consulting Provider Provider Group Practice Provider X

\* 'Consulting Provider' cannot be set to Provider Group (Required)

Consulting Provider Group

If an ordering user attempts to edit an existing non-telehealth provider consult and selects the “Provider Group” option in the “Consulting Provider” field, a new required query will appear:

- ‘Consulting Provider’ cannot be set to Provider Group

This new query is not editable, so users will never be able to fulfill the requirement and submit the order in that state.

Consult to Hospitalist Routine me 07/23/25 15:27 \* Edited

Routine \* Edited

Consulting Provider Provider Provider Group X

Consulting Provider Group

Consult on-call provider Yes No

\* Reason for consultation

\* Consulting provider notified Yes No

Only add individual provider here

Add provider groups here

For non-telehealth provider consults, users should only enter a provider group in the “Consulting Provider Group” query.

Only individual providers should be added to the “Consulting Provider” field.